

THE DELIGNE-LUSZTIG CURVE ASSOCIATED TO THE SUZUKI GROUP

FERNANDO TORRES

ABSTRACT. We give a characterization of the Deligne-Lusztig curve associated to the Suzuki group $Sz(q)$ based on the genus and the number of $\mathbb{F}_q$ -rational points of the curve.

§0. Throughout this paper by a curve we mean a projective, geometrically irreducible, and non-singular algebraic curve defined over the finite field $\mathbb{F}_q$ with q elements.

Let $N_q(g)$ denote the maximum number of $\mathbb{F}_q$ -rational points that a curve of genus g can have. The number $N_q(g)$ is bounded from above by the Hasse-Weil bound. A curve X of genus g is said to be *optimal* over $\mathbb{F}_q$, if its number of $\mathbb{F}_q$ -rational points $\#X(\mathbb{F}_q)$ is equal to $N_q(g)$. Optimal curves are very useful for applications to coding theory via Goppa construction [Go].

Besides *maximal curves* over $\mathbb{F}_q$, that is, curves whose number of $\mathbb{F}_q$ -rational points attains the Hasse-Weil bound and some curves of small genus, the only known examples of optimal curves are the Deligne-Lusztig curves associated to the Suzuki group $Sz(q)$ and to the Ree group $R(q)$ [De-Lu, §11], [Han]. Arithmetical and geometrical properties of maximal curves were studied in [R-Sti], [I], [Geer-VI] (see the references therein), [FTo] and [FGTo]. The other mentioned optimal curves were studied in [Han-Sti], [Han], [Pe] and [Han-Pe]. Hansen and Pedersen [Han-Pe, Thm. 1] stated the uniqueness, up to $\mathbb{F}_q$ -isomorphism, of the curve corresponding to $R(q)$ based on the genus, the number of $\mathbb{F}_q$ -rational points, and the group of $\mathbb{F}_q$ -automorphisms of the curve. They observed a similar result for the curve corresponding to $Sz(q)$ (cf. [Han-Pe, p.100]) as a consequence of its uniqueness up to $\mathbb{F}_q$ -isomorphism (cf. [He]). Hence, by [Han-Sti], the curve under study in this paper is $\mathbb{F}_q$ -isomorphic to the plane curve given by

$$y^q - y = x^{q_0}(x^q - x),$$

where $q_0 = 2^s$ and $q = 2q_0$. Its genus is $q_0(q - 1)$ and its number of $\mathbb{F}_q$ -rational points is $q^2 + 1$ (loc. cit.).

The paper was partially written while the author was visiting the University of Essen supported by a grant from the Graduiertenkolleg "Theoretische und experimentelle Methoden der Reinen Mathematik". The results of this paper were announced in [To].

In this note, we will be mainly concerned with the uniqueness, up to $\mathbb{F}_q$ -isomorphism, of the Deligne-Lusztig curve associated to $Sz(q)$ based on the genus and the number of $\mathbb{F}_q$ -rational points of the curve only. Our main result is the following

Theorem 1. *For $s \in \mathbb{N}$, set $q_0 := 2^{2s}$ and $q := 2q_0$. Let X be a curve over $\mathbb{F}_q$ of genus g satisfying the following hypotheses:*

1. $g = q_0(q - 1)$;
2. $\#X(\mathbb{F}_q) = q^2 + 1$.

Then X is $\mathbb{F}_q$ -isomorphic to the Deligne-Lusztig curve associated to the Suzuki group $Sz(q)$.

We also state an analogous result for the Hermitian curve over $\mathbb{F}_q$, that is, the curve defined by $y^{\sqrt{q}} + y = x^{\sqrt{q}+1}$, with q a square. This is a maximal curves over $\mathbb{F}_q$ whose genus is the biggest genus that a maximal curve over $\mathbb{F}_q$ can have, namely $\sqrt{q}(\sqrt{q} - 1)/2$ (cf. [Ih]). Several characterizations of Hermitian curves have already been given, see e.g. [HSTV] (and the references therein), [R-Sti] and [FTo]. We remark that a Hermitian curve is a Deligne-Lusztig curve associated to a projective special linear group (cf. [Han]). It is unique up to $\mathbb{F}_q$ (cf. [Han-Pe, p.100]) because of its uniqueness up to $\overline{\mathbb{F}_q}$ (cf. [Sti]). We also remark that Theorem 2 below follows from the results in [FTo] (here, we give a new proof).

Theorem 2. *Let X be a curve over $\mathbb{F}_q$ of genus g satisfying the following hypotheses:*

1. $g > (\sqrt{q} - 1)^2/4$;
2. $\#X(\mathbb{F}_q) = q + 2g\sqrt{q} + 1$ (i.e. X is maximal over $\mathbb{F}_q$).

Then X is $\mathbb{F}_q$ -isomorphic to the Hermitian curve over $\mathbb{F}_q$. In particular $g = \sqrt{q}(\sqrt{q} - 1)/2$.

To prove these theorems we use geometrical properties of the Zeta Function of curves [FGTo, §1] and a particular case of the theory of Frobenius orders associated to linear system of curves which was developed by Stöhr and Voloch [St-V] for an improvement of the Hasse-Weil bound.

It is a pleasure to thank: R. Pellikaan and H. Stichtenoth for useful conversations; A. Cossidente for having let me include his observation in this paper (see Appendix). There he points out a relation between the Deligne-Lusztig curve associated to the Suzuki group and the Suzuki-Tits ovoid. I also thank Prof. J.F. Voloch for his interest in this work.

§1. A lemma on non-gaps. Let X be a curve satisfying the hypotheses of Theorem 1, and $\text{Fr}_{\mathcal{J}}$ the Frobenius endomorphism (relative to $\mathbb{F}_q$) of the Jacobian $\mathcal{J} = \mathcal{J}_X$ of X . Then, by Serre-Weil's explicit formulae (cf. [Se], [Han]),

$$h(t) = (t^2 + 2q_0t + q)^g$$

is the characteristic polynomial of $\text{Fr}_{\mathcal{J}}$. Furthermore,

$$\text{Fr}_{\mathcal{J}}^2 + 2q_0\text{Fr}_{\mathcal{J}} + q^2I = 0 \quad \text{on } \mathcal{J},$$

due to the semi-simplicity of $\text{Fr}_{\mathcal{J}}$ and the fact that the representation of endomorphisms of $\mathcal{J}$ on the Tate module is faithful (cf. [Ta, Thm. 2], [L, VI, §3]).

Let $P_0 \in X(\mathbb{F}_q)$ and consider the natural map $f = f^{P_0} : X \rightarrow \mathcal{J}$ giving by $P \mapsto [P - P_0]$. By using $f \circ \text{Fr}_X = \text{Fr}_{\mathcal{J}} \circ f$, where Fr_X stands for the Frobenius morphism on X relative to $\mathbb{F}_q$, we then have

$$(1) \quad \text{Fr}_X^2(P) + 2q_0\text{Fr}_X(P) + qP \sim (q + 2q_0 + 1)P_0, \quad \text{for each } P \in X.$$

It turns out that X is equipped with the linear system

$$\mathcal{D} = g_{q+2q_0+1}^r := |(q + 2q_0 + 1)P_0|,$$

which is independent of $P_0 \in X(\mathbb{F}_q)$ by (1).

Lemma 1. *Let X be a curve satisfying the hypotheses of Theorem 1. For each $Q \in X(\mathbb{F}_q)$, the first positive non-gap at Q is q .*

Proof. Let $x : X \rightarrow \mathbb{P}^1(\bar{\mathbb{F}}_q)$ be such that $\text{div}_{\infty}(x) = mQ$, m being the first positive non-gap at Q . We show firstly that q is a non-gap at a generic point of X ([FGTo, proof of Prop. 1.8(i)]). Applying (as in [Har, IV, Ex. 2.6]) Fr_X to the equivalence in (1) we obtain

$$\text{Fr}_X^3(P) + (2q_0 - 1)\text{Fr}_X^2(P) + (q - 2q_0)\text{Fr}_X(P) \sim qP.$$

Consequently, if $\text{Fr}_X^i(P) \neq P$, $i = 1, 2, 3$, q is a non-gap at P . In particular

$$m \leq m_1(P) \leq q,$$

where P is a non-Weierstrass point and $m_1(P)$ is the first positive non-gap at P . On the other hand, $\#X(\mathbb{F}_q) \leq 1 + qm$ (see e.g. [Le, Thm. 1(b)]) and so $m \geq q$. Now the result follows. $\square$

§2. The linear system $\mathcal{D}$. Let X be a curve of genus g satisfying the hypothesis of the theorem. We are going to apply [St-V] to the linear system $\mathcal{D}$ introduced in §1. The key property of $\mathcal{D}$ will be the fact that $\text{Fr}_X(P)$ belongs to the tangent line at P for a generic P (see Claim 3(1)).

For $P \in X$, let $0 = j_0(P) < j_1(P) < \dots < j_r(P)$ be the $(\mathcal{D}, P)$ -orders ([St-V, §1]) and $m_i(P)$ the i th non-gap at P ($m_0(P) := 0$). If $P \in X(\mathbb{F}_q)$, from (1) we have $m_r(P) = q + 2q_0 + 1$ and

$$(2) \quad j_i(P) = m_r(P) - m_{r-i}(P) \quad \text{for } i = 0, \dots, r.$$

Then by Lemma 1 and (2),

$$(3) \quad j_r(P) = q + 2q_0 + 1, \quad j_{r-1}(P) = 2q_0 + 1 \quad \text{for } P \in X(\mathbb{F}_q).$$

Let $\epsilon_0 = 0 < \epsilon_1 < \dots < \epsilon_r$ be the orders of $\mathcal{D}$ (i.e. the $(\mathcal{D}, P)$ -orders for P generic). From (1) we see that $1, 2q_0, q$ are orders for $\mathcal{D}$. Thus $r \geq 3$ and $\epsilon_1 = 1$. Therefore

$$(4) \quad \epsilon_r = q \quad \text{and} \quad 2q_0 \leq \epsilon_{r-1} \leq 1 + 2q_0,$$

because $\epsilon_i \leq j_i(P)$ ([St-V, p.5]). Let $\nu_0 = 0 < \nu_1 < \dots < \nu_{r-1}$ be the $\mathbb{F}_q$ -Frobenius orders of $\mathcal{D}$ (which is obtained from the order sequence by deleting an specific order; cf. [St-V, Prop. 2.1]). These numbers satisfy [St-V, Corollary 2.6]

$$(5) \quad \nu_i \leq j_{i+1}(P) - j_1(P) \quad \text{whenever} \quad P \in X(\mathbb{F}_q).$$

From (1) we have that $\text{Fr}_X(P)$ belongs to the osculating hyperplane at P provided that $P \notin X(\mathbb{F}_q)$. Consequently $\nu_{r-1} = \epsilon_r = q$.

Claim 1. $\epsilon_{r-1} = 2q_0$.

Proof. Suppose that $\epsilon_{r-1} > 2q_0$. Then, by (4), $\epsilon_{r-2} = 2q_0$ and $\epsilon_{r-1} = 2q_0 + 1$. Moreover, by (5) and (3), $\nu_{r-2} \leq 2q_0 = \epsilon_{r-2}$. Hence the $\mathbb{F}_q$ -Frobenius orders of $\mathcal{D}$ would be $\epsilon_0, \epsilon_1, \dots, \epsilon_{r-2}$, and $\epsilon_r = q$.

Associated to $\mathcal{D}$ there exists an effective divisor S on X such that

$$\deg(S) = \sum_{i=0}^{r-1} \nu_i(2g-2) + (q+r)(q+2q_0+1).$$

For $P \in X(\mathbb{F}_q)$, we have

$$(6) \quad v_P(S) \geq \sum_{i=1}^r (j_i(P) - \nu_{i-1}) \geq (r-1)j_1(P) + 1 + 2q_0 \geq r + 2q_0,$$

as follows from [St-V, Prop. 2.4, Corollary 2.6], (3) and $\nu_{r-1} = q$. Thus $\deg(S) \geq (r+2q_0)\#X(\mathbb{F}_q)$, and from the formula for $\deg(S)$, $2g-2 = (2q_0-2)(q+2q_0+1)$, and $\#X(\mathbb{F}_q) = q^2+1 = (q-2q_0+1)(q+2q_0+1)$ it follows that

$$\sum_{i=1}^{r-2} \nu_i = \sum_{i=1}^{r-2} \epsilon_i \geq (r-1)q_0.$$

Now, as $\epsilon_i + \epsilon_j \leq \epsilon_{i+j}$ for $i+j \leq r$ [E, Thm. 1], then we would have

$$(r-1)\epsilon_{r-2} \geq 2 \sum_{i=0}^{r-2} \epsilon_i,$$

and hence $\epsilon_i + \epsilon_{r-2-i} = \epsilon_{r-2}$ for $i = 0, \dots, r-2$. In particular, $\epsilon_{r-3} = 2q_0 - 1$ and by the p -adic criterion (cf. [St-V, Corollary 1.9] we would have $\epsilon_i = i$ for $i = 0, 1, \dots, r-3$. These facts imply $r = 2q_0 + 2$.

Finally, we are going to see that this is a contradiction via Castelnuovo's genus bound for curves in projective spaces. We notice that $\nu_{r-2} = 2q_0$, (3) and (5) imply $j_1(P) = 1$ for $P \in X(\mathbb{F}_q)$. Then, by (2), $m_{r-1}(P) = q + 2q_0$ and hence $\mathcal{D}$ is simple.

Therefore Castelnuovo's formula (see [C], [ACGH, p. 116], [Ra, Corollary 2.8]) applied to $\mathcal{D}$ implies

$$2g = 2q_0(q-1) \leq \frac{(q+2q_0-(r-1)/2)^2}{r-1}.$$

For $r = 2q_0 + 2$ this gives $2q_0(q-1) < (q+q_0)^2/2q_0 = q_0q + q/2 + q_0/2$, a contradiction. $\square$

Claim 2. *There exists $P_1 \in X(\mathbb{F}_q)$ such that*

$$\begin{cases} j_1(P_1) = 1 \\ j_i(P_1) = \nu_{i-1} + 1 \quad \text{if } i = 2, \dots, r-1. \end{cases}$$

Proof. By (6), it is enough to show that there exists $P_1 \in X(\mathbb{F}_q)$ such that $v_{P_1}(S) = r + 2q_0$. Suppose that $v_P(S) \geq r + 2q_0 + 1$ for each $P \in X(\mathbb{F}_q)$. Then from the formula for $\deg(S)$ we would have that

$$\sum_{i=0}^{r-1} \nu_i \geq q + rq_0 + 1,$$

and, as $\epsilon_1 = 1$, $\nu_{r-1} = q$ and $\nu_i \leq \epsilon_{i+1}$, that

$$\sum_{i=0}^{r-1} \epsilon_i \geq rq_0 + 2.$$

By [E, Thm. 1], we then conclude that $r\epsilon_{r-1} \geq 2rq_0 + 4$, i.e. $\epsilon_{r-1} > 2rq_0$, a contradiction (cf. Claim 1). $\square$

Claim 3. 1. $\nu_1 > \epsilon_1 = 1$.

2. ϵ_2 is a power of two.

Proof. Statement (2) is consequence of the p -adic criterion [St-V, Corollary 1.9]. Suppose that $\nu_1 = 1$. Then by Lemma 1, Claim 2, (3) and (2) there would be a point $P_1 \in X$ such that the Weierstrass semigroup at P_1 would contain the semigroup $H := \langle q, q + 2q_0 - 1, q + 2q_0, q + 2q_0 + 1 \rangle$. Then $g \leq \#(\mathbb{N} \setminus H)$, a contradiction (cf. Remark below). $\square$

Remark. Let H be the semigroup defined above. We are going to show that $\tilde{g} := \#(\mathbb{N} \setminus H) = g - q_0^2/4$. To start with, notice that $L := \cup_{i=1}^{2q_0-1} L_i$ is a complete system

of residues module q , where

$$\begin{aligned}
L_i &= \{iq + i(2q_0 - 1) + j : j = 0, \dots, 2i\} \quad \text{if } 1 \leq i \leq q_0 - 1, \\
L_{q_0} &= \{q_0q + q - q_0 + j : j = 0, \dots, q_0 - 1\}, \\
L_{q_0+1} &= \{(q_0 + 1)q + 1 + j : j = 0, \dots, q_0 - 1\}, \\
L_{q_0+i} &= \{(q_0 + i)q + (2i - 3)q_0 + i - 1 + j : j = 0, \dots, q_0 - 2i + 1\} \cup \\
&\quad \{(q_0 + i)q + (2i - 2)q_0 + i + j : j = 0, \dots, q_0 - 1\} \quad \text{if } 2 \leq i \leq q_0/2, \\
L_{3q_0/2+i} &= \{(3q_0/2 + i)q + (q_0/2 + i - 1)(2q_0 - 1) + q_0 + 2i - 1 + j : \\
&\quad j = 0, \dots, q_0 - 2i - 1\} \quad \text{if } 1 \leq i \leq q_0/2 - 1.
\end{aligned}$$

Moreover, for each $l \in L$, $l \in H$ and $l - q \notin H$. Hence $\tilde{g}$ can be computed by summing up the coefficients of q from the above list (see e.g. [Sel, Thm. p.3]), i.e.

$$\begin{aligned}
\tilde{g} &= \sum_{i=1}^{q_0-1} i(2i+1) + q_0^2 + (q_0+1)q_0 + \sum_{i=2}^{q_0/2} (q_0+i)(2q_0-2i+2) + \\
&\quad \sum_{i=1}^{q_0/2-1} (3q_0/2+i)(q_0-2i) = q_0(q-1) - q_0^2/4.
\end{aligned}$$

In the remaining part of this section let $P_1 \in X(\mathbb{F}_q)$ be a point satisfying Claim 2. We set $m_i := m_i(P_1)$ and denote by v the valuation at P_1 .

By Claim 3(1) we have $\nu_i = \epsilon_{i+1}$ for $i = 1, \dots, r-1$. Therefore from (2), (3) and Claim 2,

$$(7) \quad \begin{cases} m_i = 2q_0 + q - \epsilon_{r-i} & \text{if } i = 1, \dots, r-2 \\ m_{r-1} = 2q_0 + q, \quad m_r = 1 + 2q_0 + q. \end{cases}$$

Let $x, y_2, \dots, y_r \in \mathbb{F}_q(X)$ be such that $\text{div}_\infty(x) = m_1 P_1$, and $\text{div}_\infty(y_i) = m_i P_1$ for $i = 2, \dots, r$. Let denote by $D^{(i)}$ the i th Hasse derivative defined by the separating variable x . The fact that $\nu_1 > 1$ means that the following matrix has rank two (cf. [St-V, proof of Prop. 2.1])

$$\begin{pmatrix} 1 & x^q & y_2^q & \dots & y_r^q \\ 1 & x & y_2 & \dots & y_r \\ 0 & 1 & D^{(1)}y_2 & \dots & D^{(1)}y_r \end{pmatrix}.$$

In particular,

$$(8) \quad y_i^q - y_i = D^{(1)}y_i(x^q - x) \quad \text{for } i = 2, \dots, r.$$

Claim 4. 1. For $P \in X(\mathbb{F}_q)$, the divisor $(2g-2)P$ is canonical, i.e. the Weierstrass semigroup at P is symmetric.

2. Let m be a non-gap at $P_1 \in X(\mathbb{F}_q)$. If $m < q + 2q_0$, then $m \leq q + q_0$.

3. For $i = 2, \dots, r$ there exists $g_i \in \mathbb{F}_q(X)$ such that $D^{(1)}y_i = g_i^{\epsilon_2}$. Furthermore, $\text{div}_\infty(g_i) = \frac{qm_i - q^2}{\epsilon_2} P_1$.

Proof. (1) Since $2g-2 = (2q_0-2)(q+2q_0+1)$, by (1) we can assume $P = P_1$. Now the case $i = r$ of Eqs. (8) implies $v(dx) = 2g-2$ and we are done.

(2) The numbers $q, q + 2q_0$ and $q + 2q_0 + 1$ are non-gaps at P_1 (cf. (7)). Then the numbers

$$(2q_0 - 2)q + q - 4q_0 + j \quad j = 0, \dots, q_0 - 2$$

are also non-gaps at P_1 . Therefore from the symmetry of the Weierstrass semigroup at P_1 ,

$$q + q_0 + 1 + j \quad j = 0, \dots, q_0 - 2$$

are gaps at P_1 . Now the proof follows.

(3) Set $f_i := D^{(1)}y_i$. By the product rule applied to (8), $D^{(j)}y_i = (x^q - x)D^{(j)}f_i + D^{(j-1)}f_i$ for $1 \leq j < q$. Then, $D^{(j)}f_i = 0$ for $1 \leq j < \epsilon_2$, because the matrices

$$\begin{pmatrix} 1 & x & y_2 & \dots & y_r \\ 0 & 1 & D^{(1)}y_2 & \dots & D^{(1)}y_r \\ 0 & 0 & D^{(j)}y_2 & \dots & D^{(j)}y_r \end{pmatrix}, \quad 2 \leq j < \epsilon_2$$

have rank two. Therefore, as ϵ_2 is a power of 2 (cf. Claim 3(2)), by [Ha-Sch, Satz 10], $f_i = g_i^{\epsilon_2}$ for some $g_i \in \mathbb{F}_q(X)$.

Finally, from the proof of item (1) we have that $x - x(P)$ is a local parameter at P if $P \neq P_1$. Then, by the election of the y_i 's, g_i has no pole but in P_1 , and from (8), $v(g_i) = -(qm_i - q^2)/\epsilon_2$. $\square$

Claim 5. $r = 4$ and $\epsilon_2 = q_0$.

Proof. We know that $r \geq 3$. We claim that $r \geq 4$; in fact, if $r = 3$ we would have $\epsilon_2 = 2q_0$, $m_1 = q$, $m_2 = q + 2q_0$, $m_3 = q + 2q_0 + 1$, and hence $v(g_2) = -q$ (g_2 being as in Claim 4(3)). Therefore, after some $\mathbb{F}_q$ -linear transformations, the case $i = 2$ of (8) reads

$$y_2^q - y_2 = x^{2q_0}(x^q - x).$$

Now the function $z := y_2^{q_0} - x^{q_0+1}$ satisfies $z^q - z = x^{q_0}(x^q - x)$ and we find that $q_0 + q$ is a non-gap at P_1 (cf. [Han-Sti, Lemma 1.8]). This contradiction eliminates the case $r = 3$.

Let $r \geq 4$ and $2 \leq i \leq r$. By Claim 4(3), $(qm_i - q^2)/\epsilon_2$ is a non-gap at P_1 , and since $(qm_i - q^2)/\epsilon_2 \geq m_{i-1} \geq q$, from (7) we have

$$2q_0 \geq \epsilon_2 + \epsilon_{r-i} \quad \text{for } i = 2, \dots, r-2.$$

In particular, $\epsilon_2 \leq q_0$. On the other hand, by Claim 4(2) we must have $m_{r-2} \leq q + q_0$ and so, by (7), we find that $\epsilon_2 \geq q_0$, i.e. $\epsilon_2 = q_0$.

Finally we show that $r = 4$. $\epsilon_2 = q_0$ implies $\epsilon_{r-2} \leq q_0$. Since $m_2 \leq q + q_0$ (cf. Claim 4(2)), by (7), we have $\epsilon_{r-2} \geq q_0$. Therefore $\epsilon_{r-2} = q_0 = \epsilon_2$, i.e. $r = 4$. $\square$

§3. Proof of Theorem 1. Let $P_1 \in X(\mathbb{F}_q)$ be as above. By (8), Claim 4(3) and Claim 5 we have the following equation

$$y_2^q - y_2 = g_2^{q_0}(x^q - x),$$

where g_2 has no pole except at P_1 . Moreover from (7) we have that $m_2 = q_0 + q$ and hence that $v(g_2) = -q$ (cf. 4(3)). Thus $g_2 = ax + b$ with $a, b \in \mathbb{F}_q$, $a \neq 0$, and after some $\mathbb{F}_q$ -linear transformations we obtain Theorem 1.

Remarks. (1) From the above computations we conclude that the Deligne-Lusztig curve associated to the Suzuki group (as in §0) is equipped with a complete and simple base-point-free linear system of dimension 4 and degree $q + 2q_0 + 1$, namely $\mathcal{D} = |(q + 2q_0 + 1)P_0|$, $P_0 \in X(\mathbb{F}_q)$. Such a linear system is an $\mathbb{F}_q$ -invariant. The orders of $\mathcal{D}$ (resp. the $\mathbb{F}_q$ -Frobenius orders) are $0, 1, q_0, 2q_0$ and q (resp. $0, q_0, 2q_0$ and q).

(2) There exists $P_1 \in X(\mathbb{F}_q)$ such that the $(\mathcal{D}, P_1)$ -orders are $0, 1, q_0 + 1, 2q_0 + 1$ and $q + 2q_0 + 1$ (cf. Claim 2, (2) and (3)). Now we show that the above sequence is, in fact, the $(\mathcal{D}, P)$ -orders for each $P \in X(\mathbb{F}_q)$. To see this, notice that

$$\deg(S) = (3q_0 + q)(2g - 2) + (q + 4)(q + 2q_0 + 1) = (4 + 2q_0)\#X(\mathbb{F}_q).$$

Let $P \in X(\mathbb{F}_q)$. From (6), we conclude that $v_P(S) = \sum_{i=1}^4 (j_i(P) - \nu_{i-1}) = 4 + 2q_0$ and hence, by (5), that $j_1(P) = 1$, $j_2(P) = q_0 + 1$, $j_3(P) = 2q_0 + 1$, and $j_4(P) = q + 2q_0 + 1$.

(3) Then, by (2), the Weierstrass semigroup $H(P)$ at $P \in X(\mathbb{F}_q)$ contains the semigroup $H := \langle q, q + q_0, q + 2q_0, q + 2q_0 + 1 \rangle$. Indeed, $H(P) = H$ since $\#(\mathbb{N} \setminus H) = g = q_0(q - 1)$ (this can be proved as in the remark after Claim 3; see also [Han-Sti, Appendix]).

(4) Let R be the ramification divisor of $\mathcal{D}$ [St-V, §1]. Its support is the set of $\mathcal{D}$ -Weierstrass points of X and

$$\deg(R) = \sum_{i=0}^4 \epsilon_i(2g - 2) + 5(q + 2q_0 + 1).$$

If $P \in X(\mathbb{F}_q)$, $v_P(R) = 2q_0 + 3$, as follows from items (1), (2) and [St-V, Thm. 1.5]. We find that $\deg(R) = (2q_0 + 3)\#X(\mathbb{F}_q)$ and consequently the set of $\mathcal{D}$ -Weierstrass points of X is equal to $X(\mathbb{F}_q)$. In particular, the $(\mathcal{D}, P)$ -orders for $P \notin X(\mathbb{F}_q)$ are $0, 1, q_0, 2q_0$ and q .

(5) We can use the above computations to obtain information on orders for the canonical morphism. By using the fact that $(2q_0 - 2)\mathcal{D}$ is canonical (cf. Claim 4(1)) and item (4), we see that the set

$$\{a + q_0b + 2q_0c + qd : a + b + c + d \leq 2q_0 - 2\}$$

is contained in the set of orders (for the canonical morphism) at non-rational points. (By considering certain rational functions of the curve, similar computations were obtained in [G-Sti, §4].)

(6) Finally, we remark that X is non-classical for the canonical morphism. We state two proofs of this fact (cf. loc. cit. and [FGTo, Prop. 1.8]): by item (5), the number q_0q is an order and since $q_0q > g - 1$ the curve is non-classical. On the other hand, from the proof of Lemma 1, q is a non-gap at a generic point P . Since $g \geq q$, this also implies the non-classicality of X .

§4. Proof of Theorem 2. The proof will be similar to the proof of Theorem 1. Let X be a curve satisfying the hypotheses of Theorem 2.

It is well known that the characteristic polynomial of the Frobenius endomorphism of the Jacobian of a maximal curve over $\mathbb{F}_q$ of genus g is $h(t) = (t + \sqrt{q})^{2g}$. Then, as in §1, X is equipped with the linear system $\mathcal{D} := g_{\sqrt{q}+1}^r = |(\sqrt{q} + 1)P_0|$ with $P_0 \in X(\mathbb{F}_q)$. Here for each $P \in X$,

$$(9) \quad \sqrt{q}P + \text{Fr}_X(P) \sim (\sqrt{q} + 1)P_0.$$

As in §1, let $j_0(P) = 0 < \dots < j_r(P)$ be the $(\mathcal{D}, P)$ -orders at P , $\epsilon_0 = 0, \dots, \epsilon_r$ (resp. $\nu_0 = 0, \dots, \nu_{r-1}$) the orders of $\mathcal{D}$ (resp. the $\mathbb{F}_q$ -Frobenius orders of $\mathcal{D}$). By (9), $\nu_{r-1} = \epsilon_r$, $j_r(P) = m_r(P) = \sqrt{q} + 1$ for each $P \in X(\mathbb{F}_q)$, and 1 and $\sqrt{q}$ are $(\mathcal{D}, P)$ -orders for each $P \notin X(\mathbb{F}_q)$. Then $j_1(P) = 1$ and $\sqrt{q} \leq j_r(P) \leq \sqrt{q} + 1$ at $P \notin X(\mathbb{F}_q)$. In this case we have $j_r(P) = \sqrt{q}$: in fact, if $j_r(P) = \sqrt{q} + 1$, by (9), we would have $(\sqrt{q} + 1)P \sim \sqrt{q}P + \text{Fr}_X(P)$, i.e. $P \sim \text{Fr}_X(P)$ and so $g = 0$, a contradiction. Therefore $\nu_{r-1} = \epsilon_r = \sqrt{q}$. Now, the invariants of $\mathcal{D}$ also satisfy properties (2) and (5) above. The case $i = r - 1$ of (5) gives $j_1(P) = 1$ for each $P \in X(\mathbb{F}_q)$. Hence by (2) we find that $m_{r-1}(P) = \sqrt{q}$ for each $P \in X(\mathbb{F}_q)$. It follows that $\mathcal{D}$ is simple. Now by using the hypothesis on g and Castelnuovo's genus bound we find that $r = 2$ (cf. [FTo, Claim 1]).

Then, by using the ramification divisor of $\mathcal{D}$, we have that $g = \sqrt{q}(\sqrt{q} - 1)/2$ (see [FTo, §3]). Fix $P_0 \in X(\mathbb{F}_q)$, $x, y \in \mathbb{F}_q(X)$ with $\text{div}_\infty(x) = \sqrt{q}P_0$, and $\text{div}_\infty(y) = (1 + \sqrt{q})P_0$. Then $H(P_0) = \langle \sqrt{q}, \sqrt{q} + 1 \rangle$ and consequently $\text{div}(dx) = (2g - 2)P_0(*)$, because $H(P_0)$ is symmetric.

By $\nu_1 = \sqrt{q}$ we have an equation of type

$$(10) \quad y^q - y = f(x^q - x),$$

with $f := D^1y$ (derivation with respect to x), and by $\epsilon_2 = \sqrt{q}$ we have the following rank 2 matrices

$$\begin{pmatrix} 1 & x & y \\ 0 & 1 & D^{(1)}y \\ 0 & 0 & D^{(j)}y \end{pmatrix}, \quad 2 \leq j < \epsilon_2 = \sqrt{q}.$$

By (*) and (10), $\text{div}_\infty(f) = qP_0$. Then $D^{(j)}y = 0$ for $2 \leq j < \sqrt{q}$ and from (10) we have that $D^{(j)}f = 0$ for $1 \leq j < \sqrt{q}$. Then by [Ha-Sch, Satz 10], there exists $f_1 \in \mathbb{F}_q(X)$ such that $f = f_1^{\sqrt{q}}$. Therefore $f_1 = ax + b$ with $a, b \in \mathbb{F}_q$, $a \neq 0$, and after some $\mathbb{F}_q$ -linear transformations we obtain an equation of type

$$y_1^{\sqrt{q}} + y_1 - x_1^{\sqrt{q}+1} = (y_1^{\sqrt{q}} + y_1 - x_1^{\sqrt{q}+1})^{\sqrt{q}},$$

with $x_1, y_1 \in \mathbb{F}_q(X)$. Now the proof follows.

Remark. Similar computations to those of the Remarks in §3 can be obtained in this case. In particular, using the divisor $\mathcal{D}$, one can show that Hermitian curves are non-classical curves (see also [G-Via]). We mention that the first examples of non-classical curves were obtained from some Hermitian curves (see [Sch]).

Appendix: On the Suzuki-Tits ovoid

For $s \in \mathbb{N}$, let $q_0 := 2^s$ and $q := 2q_0$. It is well known that the Suzuki-Tits ovoid $\mathcal{O}$ can be represented in $\mathbb{P}^4(\mathbb{F}_q)$ as

$$\mathcal{O} = \{(1 : a : b : f(a, b) : af(a, b) + b^2) : a, b \in \mathbb{F}_q\} \cup \{(0 : 0 : 0 : 0 : 1)\},$$

where $f(a, b) := a^{2q_0+1} + b^{2q_0}$ (see [Tits], [Pen-W, p.3])

Let X be the Deligne-Lusztig curve associated to $Sz(q)$ and $\mathcal{D} = |(q + 2q_0 + 1)P_0|$, $P_0 \in X(\mathbb{F}_q)$ (see §1). By Remark §3(3), we can associate to $\mathcal{D}$ a morphism $\pi = (1 : x : y : z : w)$ whose coordinates satisfy $\text{div}_\infty(x) = qP_0$, $\text{div}_\infty(y) = (q + q_0)P_0$, $\text{div}_\infty(z) = (q + 2q_0)P_0$ and $\text{div}_\infty(w) = q + 2q_0 + 1$.

Claim. (*A. Cossidente*) $\mathcal{O} = \pi(X(\mathbb{F}_q))$.

Proof. We have $\pi(P_0) = (0 : 0 : 0 : 0 : 1)$; we can choose x and y satisfying the plane equation (given in §0) of the curve, $z := x^{2q_0+1} + y^{2q_0}$, and $w := xy^{2q_0} + z^{2q_0} = xy^{2q_0} + x^{2q+2q_0} + y^{2q}$ (cf. [Han-Sti, S1.7]). For $P \in X(\mathbb{F}_q) \setminus \{P_0\}$ set $a := x(P)$, $b := y(P)$, and $f(a, b) := z(a, b)$. Then $w(a, b) = af(a, b) + b^2$ and we are done. $\square$

Remark. The morphism π is an embedding. Indeed, since $j_1(P) = 1$ for each P (cf. Remarks §3(2)(4)), it is enough to see that π is injective. By (1), the points P where π could not be injective satisfy: $P \notin X(\mathbb{F}_q)$, $\text{Fr}_X^3(P) = P$ or $\text{Fr}_X^2(P) = P$. Now from the Zeta function of X one sees that $\#X(\mathbb{F}_{q^3}) = \#X(\mathbb{F}_{q^2}) = \#X(\mathbb{F}_q)$, and the remark follows.

REFERENCES

- [ACGH] Arbarello, E.; Cornalba, M.; Griffiths, P.A. and Harris, J.: Geometry of algebraic curves, Vol I, Springer Verlag, New York 1985.
- [C] Castelnuovo, G.: Ricerche di geometria sulle curve algebriche, Atti. R. Acad. Sci. Torino **24**, 196–223 (1889).

- [De-Lu] Deligne, P.; Lusztig: Representations of reductive groups over finite fields, *Ann. of Math.* **103**, 103–161 (1976).
- [E] Esteves, E.: A geometric proof of an inequality of order sequences, *Comm. Algebra* **21**(1), 231–238 (1993).
- [FTo] Fuhrmann, R.; Torres, F.: The genus of curves over finite fields with many rational points, *Manuscripta Math.* **89**, 103–106 (1996).
- [FGTo] Fuhrmann, R.; Garcia, A.; Torres, F.: On maximal curves, to appear in JNT.
- [G-Sti] Garcia, A.; Stichtenoth, H.: Elementary abelian p -extensions of algebraic functions fields, *Manuscripta Math.* **72**, 67–79 (1991).
- [G-Via] Garcia, A.; Viana, P.: Weierstrass points on certain non-classical curves, *Arch. Math.* **46**, 315–322 (1986).
- [Geer-Vl] van der Geer, G.; van der Vlugt, M.: How to construct curves over finite fields with many points, *alg-geom/ 9511005*.
- [Go] Goppa, V.D.: Algebraic-Geometric codes, *Math. USSR-Izv.* **21** (1), 75–91 (1983).
- [Han] Hansen, J.P.: Deligne-Lusztig varieties and group codes, *Lect. Notes Math.* **1518**, 63–81 (1992).
- [Han-Pe] Hansen, J.P.; Pedersen, J.P.: Automorphism group of Ree type, Deligne-Lusztig curves and function fields, *J. reine angew. Math.* **440**, 99–109 (1993).
- [Han-Sti] Hansen, J.P.; Stichtenoth, H.: Group codes on certain algebraic curves with many rational points, *AAECC* **1**, 67–77 (1990).
- [Har] Hartshorne, R.: *Algebraic Geometry*, Grad. Texts in Math. **52** Springer-Verlag 1977.
- [Ha-Sch] Hasse, H.; Schmidt, F.K.: Noch eine Begründung der Theorie der höheren Differentialquotienten in einem algebraischen Funktionenkörper einer Unbestimmten, *J. reine angew. Math.* **177**, 215–237 (1937).
- [He] Henn H.W.: Funktionenkörper mit großer Automorphismengruppe, *J. reine angew. Math.* **302**, 96–115 (1978).
- [HSTV] Hirschfeld, J.W.P.; Storme, L.; Thas, J.A. and Voloch, J.F.: A characterization of Hermitian curves, *J. Geom.* **41**, 72–78 (1991).
- [I] Ibukiyama, T.: On rational points of curves of genus 3 over finite fields, *Tôhoku Math. J.* **45**, 311–329 (1993).
- [Ih] Ihara, Y.: Some remarks on the number of rational points of algebraic curves over finite fields, *J. Fac. Sci. Tokio* **28**, 721–724 (1981).
- [L] Lang, S.: *Abelian Varieties*, Interscience Pub., New York 1959.
- [Le] Lewittes, J.: Places of degree one in function fields over finite fields, *J. Pure Appl. Alg.* **69**, 177–183 (1990).
- [Pe] Pedersen, J.P.: A function field related to the Ree group, *Lect. Notes Math.* **1518**, 122–131 (1992).
- [Pen-W] Penttila, T.; Williams, B.: Ovoids of parabolic spaces, preprint (1997).
- [Ra] Rathmann, J.: The uniform position principle for curves in characteristic p , *Math. Ann.* **276**, 565–579 (1987).
- [R-Sti] Rück, H.G.; Stichtenoth, H.: A characterization of Hermitian function fields over finite fields, *J. reine angew. Math.* **457**, 185–188 (1994).
- [Sch] Schmidt, F.K.: Zur arithmetischen Theorie der algebraischen Funktionen II. Allgemeine Theorie der Weierstrasspunkte, *Math. Z.* **45**, 75–96 (1939).
- [Sel] Selmer, E.S.: On the linear diophantine problem of Frobenius, *J. reine angew. Math.* **293/294**, 1–17 (1977).

- [Se] Serre, J.P.: Sur le nombre des points rationnels d'une courbe algébrique sur un corps fini, C.R. Acad. Sci. Paris Sér. I Math. **296**, 397–402 (1983).
- [Sti] Stichtenoth, H.: Über die Automorphismengruppe eines algebraischen Funktionenkörpers von Primzahlcharakteristik. Teil I: Eine Abschätzung der Ordnung der Automorphismengruppe, Arch. Math. **24**, 527–544 (1973).
- [St-V] Stöhr, K.O.; Voloch, J.F.: Weierstrass points and curves over finite fields, Proc. London Math. Soc. (3) **52**, 1–19 (1986).
- [Ta] Tate, J.: Endomorphisms of abelian varieties over finite fields, Inventiones Math. **2**, 134–144 (1966).
- [Tits] Tits, J.: Ovoïdes et groupes de Suzuki, Arch. Math. **13**, 187–198 (1962).
- [To] Torres, F.: A characterization of some optimal curves, Finite Geometry and Combinatorics, Third International Conference at Deinze, May 1997.

ICTP, MATHEMATICS SECTION, P.O. BOX 586, 34100, TRIESTE, ITALY

E-mail address: ftorres@ictp.trieste.it